

## DRUG FACTS

### Active ingredient(s)

### Purpose

#### Use(s)

temporarily relieves these symptoms associated with the common cold or flu:

- headache
- fever
- sinus pressure
- nasal congestion
- minor body aches and pains

#### Warnings

##### Do not use

- in children under 12 years of age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

##### Ask a doctor before use if you have

- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- bleeding problems
- high blood pressure
- heart or kidney disease
- asthma
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- taken a diuretic
- reached age 60 or older

##### Ask a doctor or pharmacist before use if you are

- taking any other drug containing a NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor's care for any serious condition
- taking any other product that contains pseudoephedrine or any other nasal decongestant
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

##### When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

##### Stop use and ask a doctor if

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- symptoms continue or get worse
- stomach pain or upset gets worse or lasts
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- any new symptoms appear

##### If pregnant or breast-feeding

##### Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

#### Other information

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- **do not take more than directed**
- **the smallest effective dose should be used**
- do not take longer than 7 days, unless directed by a doctor (see WARNINGS)
- adults and children 12 years of age and over:
- take 1 capsule every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 capsule, 2 capsules may be used.
- do not use more than 6 capsules in any 24-hour period unless directed by a doctor
- children under 12 years of age: do not use
- **each capsule contains:** potassium 20 mg
- store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F).

- read all warnings and directions before use. Keep carton.

***Inactive ingredients***

D&C yellow no. 10, FD&C red no. 40, fractionated coconut oil, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol